



AgraQuest, Inc.
 1530 Drew Avenue
 Davis, CA 95616
 530-750-0150
 530-750-0153 Fax
 www.agraquest.com
 info@agraquest.com

Serenade® MAX EPA Reg. No. 69592-11

SUPPLEMENTAL LABEL

(Includes uses on Citrus and Fruiting Vegetables not previously on Label Codes #40CI-001 and 40BBA-001)

Recommended Application Rates for Selected Crops – Agricultural Use

Serenade MAX has a 0-Day PreHarvest Interval for all crops contained on this label

Under moderate to severe disease pressure, for improved performance, increase rates and reduce spray intervals or use Serenade MAX in a tank mix or rotational program with other registered fungicides.

Crops	Disease	Rate lbs./acre [#]	Application Instructions
Citrus Orange Grapefruit Lemon Tangerine Tangelo Pummelo and other citrus crops	Bacterial Canker <i>Xanthomonas spp</i> Bacterial Blast <i>Peudomonas syringae</i>	1 - 3	Bacterial Canker/Bacterial Blast – Begin applications when environmental conditions are conducive to disease development. Repeat on 7 to 10 day intervals or as needed.
Fruiting Vegetables Pepper Tomato Eggplant Ground Cherry Tomatillo Okra and other fruiting vegetables	Target Spot <i>Corynespora casslicola</i>	1 - 3	Begin application soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a 5 to 7 day interval or as needed. When conditions are conducive to rapid disease development, for improved control, use Serenade in a tank mix program with copper-based bactericides at labeled rates.
	Bacterial Canker <i>Clavibacter michiganensis</i>	1 - 3	Begin applications when environmental conditions are conducive to disease development. Repeat on 7 to 10 day intervals or as needed.

[See basic label attached to container for First Aid, Precautionary Statements and additional Directions for Use]